

---

# Anatomy Of The Lower Limb

---

*Anatomy Of  
The Lower  
Limb*

*Downloaded from  
[web1.enteraskincare.com](http://web1.enteraskincare.com)  
by Lozano & Beltran*

---

## INTRODUCTION TO THE ANATOMY OF THE LOWER LIMB

---

The human lower limb is a remarkable feat of biomechanical engineering, designed to support the entire weight of the body while providing stability for standing and the explosive power needed for walking, running, and jumping.

Understanding the **anatomy of the lower limb** is essential not only for medical professionals like orthopedists and

physiotherapists but also for athletes, fitness enthusiasts, and anyone interested in how their body moves. This complex region is composed of bones, joints, muscles, nerves, blood vessels, and connective tissues, all working in perfect harmony. In this comprehensive guide, we will explore the intricate structure of the lower limb, from the hip to the toes, breaking down each component and explaining its function in a way that is both informative and easy to follow.

---

## LOWER LIMB

---

The lower limb is divided into several distinct anatomical regions, each with its own specialized function. These regions include the gluteal region (the buttock), the thigh (between the hip and the knee), the leg (specifically the portion between the knee and the ankle, often referred to as the lower leg), and the foot. The **anatomy of the lower limb** is organized to maximize both strength and mobility, allowing for a wide range of motion while maintaining the structural integrity needed to bear weight. The bones of the lower limb are thicker and stronger than those of the upper limb, reflecting their role in

weight support and locomotion. Similarly, the muscles are larger and more powerful, designed for endurance and force production.

## THE BONY FRAMEWORK: BONES OF THE LOWER LIMB

---

The skeletal foundation of the lower limb provides the rigid framework upon which all other tissues attach. It consists of 30 bones per limb, including the hip bone, femur, patella, tibia, fibula, and the bones of the foot. Each bone plays a critical role in the overall function of the limb.

## The Hip

# Bone

The hip bone, or os coxae, is a large, irregularly shaped bone that forms the pelvic girdle. It is actually composed of three fused bones: the ilium, ischium, and pubis. The hip bone articulates with the sacrum at the sacroiliac joint and with the femur at the hip joint. The acetabulum, a deep socket on the lateral surface of the hip bone, receives the head of the femur, forming the hip joint. The architecture of the hip bone is designed to transmit weight from the axial skeleton to the lower limbs.

# The

# Femur

The femur, or thigh bone, is the longest, heaviest, and strongest bone in the human body. It extends from the hip joint to the knee joint. The proximal end features a rounded head that fits into the acetabulum, a neck that is a common site for fractures in the elderly, and two prominences called the greater and lesser trochanters, which serve as attachment points for powerful muscles. The shaft of the femur is slightly bowed anteriorly, which helps distribute mechanical stress. The distal end expands into two rounded condyles that articulate with the tibia and patella to form the knee joint.

# The Patella

The patella, commonly known as the kneecap, is a large sesamoid bone embedded within the quadriceps tendon. It sits anterior to the knee joint and serves to protect the joint, increase the leverage of the quadriceps muscle, and reduce friction between the tendon and the femur. The patella is a crucial component of the **anatomy of the lower limb** for activities that involve knee extension.

# The Tibia

The tibia, or shin bone, is the larger and more medial of the two bones in the leg. It bears the majority of

the body weight transmitted from the femur. The proximal end features two flat articular surfaces called the medial and lateral tibial plateaus, which articulate with the femoral condyles. The tibial tuberosity, a bony prominence on the anterior surface, is the attachment site for the patellar ligament. The distal end of the tibia is smaller and articulates with the talus bone of the foot to form the ankle joint, with the medial malleolus forming the bony bump on the inside of the ankle.

# The Fibula

The fibula is a long, slender bone located laterally to the tibia. It does not bear

significant weight but serves as an important attachment site for muscles and ligaments. The proximal head of the fibula articulates with the lateral condyle of the tibia, while the distal end forms the lateral malleolus, the bony bump on the outside of the ankle. The fibula is an integral part of the **anatomy of the lower limb** for ankle stability and muscle function.

## Bones of the Foot

The foot is a complex structure composed of 26 bones, which are organized into three groups: the tarsals, metatarsals, and phalanges. The seven tarsal bones include the talus (which articulates with the

tibia and fibula to form the ankle), the calcaneus (the heel bone), the navicular, the cuboid, and the three cuneiforms. The five metatarsal bones form the midfoot, and the phalanges form the toes. The arch of the foot, maintained by bones and ligaments, acts as a natural shock absorber during walking and running.

---

### THE MUSCULAR SYSTEM OF THE LOWER LIMB

---

The muscles of the lower limb are among the most powerful in the body. They are organized into compartments, each with specific functions related to movement and stability. A thorough understanding of the muscular **anatomy of the lower limb** is key

to comprehending how we walk, run, jump, and maintain posture.

## Muscles of the Gluteal Region

The gluteal region contains the gluteal muscles, which are primarily responsible for hip extension, abduction, and rotation. The gluteus maximus is the largest muscle in the body and is the primary hip extensor, crucial for climbing stairs and running. The gluteus medius and gluteus minimus are important for hip abduction and pelvic stabilization during the single-leg stance phase of gait.

These muscles are often targeted in rehabilitation and fitness training due to their role in lower limb function.

## Muscles of the Thigh

The thigh muscles are divided into three compartments: the anterior, medial, and posterior.

- **Anterior compartment:**

This compartment contains the quadriceps femoris, a group of four muscles (rectus femoris, vastus lateralis, vastus medialis, and vastus intermedius) that

are the primary extensors of the knee. It also includes the sartorius, the longest muscle in the body, which assists in hip and knee flexion.

- **Medial compartment:** These muscles, collectively known as the adductors (adductor longus, adductor brevis, adductor magnus, gracilis, and pectineus), are primarily responsible for adduction of the hip, or bringing the thigh toward the midline.
- **Posterior compartment:** This compartment contains the hamstring

muscles (biceps femoris, semitendinosus, and semimembranosus), which are the primary flexors of the knee and extensors of the hip. These muscles are frequently injured in sports that involve sprinting and sudden acceleration.

## Muscles of the Leg

The leg, or lower leg, also has three compartments: anterior, lateral, and posterior.

- **Anterior**

- **compartment:**

These muscles (tibialis anterior, extensor digitorum longus, extensor hallucis longus, and fibularis tertius) are responsible for dorsiflexion of the ankle (lifting the foot upward) and extension of the toes. The tibialis anterior is a key muscle for foot clearance during the swing phase of gait.

- **Lateral**

- **compartment:**

The fibularis longus and fibularis brevis are primarily responsible for eversion of the foot (turning the sole outward) and assist in plantarflexion.

- **Posterior**

- **compartment:**

This is the largest compartment and contains the powerful plantarflexors. The superficial group includes the gastrocnemius and soleus, which form the calf and converge into the Achilles tendon. The deep group includes the popliteus, tibialis posterior, flexor digitorum longus, and flexor hallucis longus, which are involved in plantarflexion, inversion of the foot, and flexion of the toes.

# Intrinsic Muscles of the Foot

The foot also contains numerous small intrinsic muscles that control the fine movements of the toes and help maintain the arches of the foot. These include the flexor digitorum brevis, abductor hallucis, and the interossei muscles. They are essential for balance and propulsion during gait.

---

## THE VASCULAR SYSTEM: ARTERIES AND VEINS OF THE LOWER LIMB

---

The blood supply to the lower limb is critical for delivering oxygen and

nutrients to the tissues and removing waste products. The arterial system begins with the femoral artery, which is a continuation of the external iliac artery. The femoral artery travels through the thigh, giving off several branches, including the profunda femoris artery, which supplies the deep muscles of the thigh. As it passes posterior to the knee, the femoral artery becomes the popliteal artery, which then divides into the anterior and posterior tibial arteries. The anterior tibial artery supplies the anterior compartment of the leg and becomes the dorsalis pedis artery on the dorsum of the foot. The posterior tibial artery supplies the posterior compartment and gives off the

fibular artery. The venous system includes superficial veins, such as the great saphenous vein and small saphenous vein, which drain into the deep veins. The deep veins, such as the femoral and popliteal veins, accompany the arteries. The venous system relies on valves and muscle contractions to return blood to the heart, which is a key concept in the **anatomy of the lower limb** for understanding conditions like deep vein thrombosis.

---

## NERVES OF THE LOWER LIMB

---

The lower limb is innervated by branches of the lumbosacral plexus, which arises from the spinal nerves of the lumbar and sacral

regions. The major nerves include the femoral nerve and the sciatic nerve.

- **Femoral nerve:**

Arising from the lumbar plexus, the femoral nerve innervates the anterior compartment muscles of the thigh and gives sensory branches to the anterior thigh and medial leg.

- **Sciatic nerve:**

The largest nerve in the body, the sciatic nerve arises from the sacral plexus and travels through the posterior thigh. It gives off branches to the hamstring muscles and then divides into the tibial nerve and the common

fibular (peroneal) nerve at the knee. The tibial nerve innervates the posterior compartment of the leg and the sole of the foot, while the common fibular nerve innervates the anterior and lateral compartments of the leg.

Understanding the course of these nerves is essential in the clinical **anatomy of the lower limb**, as they are vulnerable to injury from trauma, compression, or prolonged pressure.

- **Obturator**

**nerve:** This nerve arises from the lumbar plexus and innervates the adductor muscles of the medial thigh.

---

## JOINTS AND ARTICULATIONS

---

The joints of the lower limb provide both stability and mobility. The major joints are the hip, knee, and ankle, each with a unique structure and function.

# The Hip Joint

The hip joint is a ball-and-socket synovial joint formed between the head of the femur and the acetab