

---

# How To Increase Blood Pressure

---

*How To Increase Blood Pressure* Downloaded from  
[web1.enteraskincare.com](http://web1.enteraskincare.com)  
by Pitts & Black

---

## UNDERSTANDING LOW BLOOD PRESSURE AND WHY RAISING IT MATTERS

---

When we talk about cardiovascular health, the conversation usually centers on high blood pressure. However, for millions of people, the opposite condition—hypotension, or chronically low blood pressure—can be just as disruptive. While a reading of 120/80 mmHg is often considered ideal, some individuals consistently measure below 90/60 mmHg, which can lead to dizziness, fatigue, fainting, and even shock in severe cases. Knowing **how to increase blood pressure** safely and effectively is essential for anyone who experiences persistent symptoms or has been diagnosed with hypotension by a healthcare provider.

Low blood pressure is not always a medical problem. In fact, many physically fit people have naturally low readings without any negative effects. However, when blood pressure drops too low and the brain and vital organs do not receive enough oxygen and nutrients, the body sends clear warning signals. These include blurred vision, nausea, lack of concentration, cold and clammy skin, rapid shallow breathing, and a feeling of lightheadedness upon standing. Before rushing to raise your numbers, it is important to understand the underlying cause, because the right approach depends on whether the low

pressure stems from dehydration, medication side effects, nutritional deficiencies, or an underlying health condition like diabetes or heart disease.

---

## THE ROLE OF HYDRATION AND FLUID VOLUME

---

### Why Dehydration Lowers Blood Pressure

One of the most straightforward and effective methods for **how to increase blood pressure** is to ensure adequate hydration. The human body is roughly 60% water, and blood volume is directly tied to fluid intake. When you are dehydrated, the total volume of blood circulating through your veins decreases, which can cause a significant drop in pressure. Even mild dehydration—losing as little as 1.5% of your body's water—can make blood pressure dip enough to trigger symptoms. The fix is simple: drink water consistently throughout the day. For active individuals or those living in hot climates, increasing intake further can make a noticeable difference.

### How Much Fluid Is Enough?

General guidelines suggest eight to ten 8-ounce glasses of water daily, but

needs vary. A practical test is to monitor the color of your urine; pale yellow indicates proper hydration, while dark yellow or amber suggests you need more fluids. In addition to plain water, clear soups, herbal teas, and high-water-content fruits like watermelon and cucumber count toward your total intake. However, avoid excessive consumption of caffeinated beverages and alcohol, as both can act as diuretics and further lower blood pressure by promoting fluid loss.

---

### DIETARY ADJUSTMENTS: SODIUM AND BEYOND

---

## The Strategic Use of Salt

Sodium has a direct and powerful effect on blood pressure. While the general population is advised to limit salt to manage hypertension, individuals with chronic low blood pressure may benefit from a moderate increase in dietary sodium. Adding a pinch of high-quality sea salt to meals, or choosing naturally salty foods like olives, pickles, and canned fish (in moderation), can help raise volume in the bloodstream. The trick is balance—too much salt can eventually lead to other health problems, including kidney strain and fluid retention in the wrong places. Therefore, anyone exploring **how to increase blood pressure** with salt should discuss it with a doctor first, especially if they have a history of heart failure or kidney disease.

## Smaller, More Frequent Meals

Postprandial hypotension is a common phenomenon in which blood pressure drops significantly after eating, particularly after large meals. This happens because digestion diverts blood flow to the stomach and intestines. A simple and effective strategy is to eat smaller, more frequent meals throughout the day rather than three large ones. This approach helps keep digestion steady and prevents sudden drops in pressure. Additionally, including complex carbohydrates and lean proteins can stabilize blood sugar and energy levels, both of which influence circulatory function.

## Foods That Naturally Support Blood Pressure

Certain foods contain nutrients that help maintain healthier blood pressure levels. For example, foods rich in vitamin B12 and folate are essential because deficiencies in these vitamins can lead to anemia, a known cause of low blood pressure. Include eggs, fortified cereals, lean meat, and leafy greens in your diet. Also, licorice root has been shown to raise blood pressure in some people, though it must be used cautiously due to potential side effects like low potassium. A more practical option is to drink beetroot juice, which contains nitrates

that may help regulate circulation. Always consider the whole dietary pattern rather than relying on a single superfood.

---

### LIFESTYLE CHANGES FOR STEADIER CIRCULATION

---

## Movement and Compression

Physical inactivity can cause blood to pool in the legs, leading to a drop in central blood pressure. Regular, gentle exercise like walking, swimming, or cycling helps the cardiovascular system become more responsive. However, avoid sudden, intense exertion that could cause a sharp drop. Compression stockings are another powerful non-medical tool. By applying graduated pressure to the legs, they reduce the amount of blood that pools in the lower extremities, helping to maintain adequate pressure in the upper body. This is particularly helpful for people who experience orthostatic hypotension—dizziness upon standing—and is a key recommendation when learning **how to increase blood pressure** naturally.

## Changing Positions Slowly

One of the simplest adjustments with immediate results is to change positions gradually. If you feel dizzy when standing up, try sitting up for a minute or two first, then stand slowly while holding onto a stable surface. This gives your autonomic nervous system time to adjust and constrict blood vessels

appropriately. Over time, this habit can prevent fainting episodes and reduce the risk of injury from falls.

## Adequate Sleep and Stress Management

Poor sleep and chronic stress can disrupt the delicate balance of the autonomic nervous system, which regulates blood pressure. Aim for seven to nine hours of quality rest per night, and incorporate relaxation techniques such as deep breathing, meditation, or gentle yoga. Avoid long periods of standing still, especially in hot environments, as this can cause blood to pool and pressure to drop. If you must stand for long periods, shifting weight from one foot to the other or crossing your legs can help pump blood back upward.

---

### MEDICAL AND PHARMACOLOGICAL OPTIONS

---

## Reviewing Current Medications

Many cases of low blood pressure are linked to prescribed medications. Diuretics, alpha blockers, beta blockers, antidepressants, and drugs for erectile dysfunction are just a few examples that can inadvertently lower pressure. Before trying any intervention, a thorough review of your medication list with a healthcare professional is essential. Sometimes, adjusting the dose or

timing—or switching to a different drug—can resolve the problem without needing additional treatments. Never stop or change a prescription on your own.

## Fludrocortisone and Midodrine

For chronic hypotension that does not respond to lifestyle and dietary changes, doctors may prescribe medications. Fludrocortisone helps the body retain sodium and fluid, thereby increasing blood volume. Midodrine works by constricting blood vessels to raise pressure. These drugs are typically reserved for cases where symptoms are debilitating and other methods have failed. They require careful monitoring because they can cause side effects like headache, supine hypertension (high blood pressure while lying down), and fluid imbalances. Understanding **how to increase blood pressure** with medical help means fully discussing risks and benefits with your doctor.

## Treating Underlying Conditions

Sometimes low blood pressure is a symptom of another problem: anemia, thyroid issues, adrenal insufficiency, diabetes, or heart valve abnormalities. In these cases, the most effective way to raise blood pressure is to treat the root cause. For example, iron supplements can correct anemia, thyroid hormone replacement can stabilize metabolism,

and insulin management can prevent diabetic autonomic neuropathy. Addressing these conditions often leads to a natural improvement in blood pressure without direct intervention.

---

### WHEN TO SEEK EMERGENCY CARE

---

While many people can manage mild hypotension with lifestyle changes, there are clear warning signs that require immediate medical attention. If low blood pressure is accompanied by chest pain, shortness of breath, high fever, severe headache, stiff neck, or loss of consciousness, call emergency services. These symptoms could indicate a serious condition such as sepsis, a heart attack, anaphylaxis, or internal bleeding. Similarly, if you notice blood in your stool or vomit, or if you feel your heart racing uncontrollably, do not attempt to self-treat. The priority is ruling out life-threatening causes before working on long-term management.

---

### PUTTING IT ALL TOGETHER: A PERSONALIZED APPROACH

---

There is no one-size-fits-all answer to **how to increase blood pressure**. The right strategy depends on your medical history, lifestyle, and the underlying reason for the low readings. Start with the least invasive methods: drink more water, increase salt intake moderately, eat small meals, and wear compression stockings. Monitor your symptoms and take note of what works. Keep a log of your blood pressure readings at different times of day, especially after eating or standing. Share this information with your healthcare provider to refine your plan.

Patience is important. Blood pressure can fluctuate from hour to hour, and

changes from diet or medication may take days or weeks to show consistent results. Never try to raise your pressure too quickly; a sudden spike can be just as dangerous as a chronic low. The goal is not to achieve a perfect number, but to feel steady, clear-headed, and energetic throughout the day.

---

## CONCLUSION

---

Low blood pressure can be a frustrating and worrisome condition, but in most cases it can be managed effectively with a combination of smart hydration, targeted nutrition, thoughtful lifestyle

changes, and medical guidance when needed. The key is to address the root cause rather than simply chasing a number on a monitor. By learning **how to increase blood pressure** safely—whether through a pinch of salt, a pair of compression socks, or a medication adjustment—you can regain control over your health and prevent the dizzying, draining symptoms that disrupt daily life. Always work closely with your doctor to find the balance that works for you, and remember that small, consistent changes often yield the most sustainable results.